

Endocrinology

Abstract

Endocrinology (from [Greek](#) [ἔνδον](#), *endon*, "within"; [κρίνω](#), *krīnō*, "to separate"; and [-λογία](#), *-logia*) is a branch of medicine dealing with disorder of the [endocrine system](#) and its specific secretions called [hormones](#), the integration of developmental events such as proliferation, growth, and differentiation (including [histogenesis](#) and [organogenesis](#)) and the coordination of [metabolism](#), [respiration](#), excretion, movement, [reproduction](#), and sensory perception depend on chemical cues, substances synthesised and secreted by specialized cells.

Endocrinology is concerned with the study of the biosynthesis, storage, chemistry, and physiological function of [hormones](#) and with the cells of the endocrine glands and tissues that secrete them.

The endocrine system consists of several glands, in different parts of the body, that secrete hormones directly into the blood rather than into a duct system. Hormones have many different functions and modes of action; one hormone may have several effects on different target organs, and, conversely, one target organ may be affected by more than one hormone.


In the original 1902 definition by Bayliss and Starling (see below), they specified that, to be classified as a hormone, a chemical must be produced by an organ, be released (in small amounts) into the blood, and be transported by the blood to a distant organ to exert its specific function. This definition holds for most "classical" hormones, but there are also [paracrine](#) mechanisms (chemical communication between cells within a tissue or organ), autocrine signals (a chemical that acts on the same cell), and [intracrine](#) signals (a chemical that acts within the same cell).^[1] A [neuroendocrine](#) signal is a "classical" hormone that is released into the blood by a neurosecretory neuron (see article on [Neuroendocrinology](#)).

Hormones act by binding to specific [receptors](#) in the target organ. As [Baulieu](#) notes, a receptor has at least two basic constituents:


- a recognition site, to which the hormone binds
- an effector site, which precipitates the modification of cellular function.^[2]

Between these is a "transduction mechanism" in which hormone binding induces allosteric modification that, in turn, produces the appropriate response.

Chemical classes of hormones



Amine hormones: norepinephrine and triiodothyronine



Steroid hormones: cortisol and vitamin D3

Griffin and Ojeda identify three different classes of hormone based on their chemical composition:^[3]

Amines

Amines, such as [norepinephrine](#), [epinephrine](#), and [dopamine](#), are derived from single amino acids, in this case tyrosine. [Thyroid](#) hormones such as 3,5,3'-triiodothyronine (T₃) and 3,5,3',5'-tetraiodothyronine (thyroxine, T₄) make up a subset of this class because they derive from the combination of two iodinated tyrosine amino acid residues.

Peptide and protein

[Peptide hormones](#) and protein hormones consist of three (in the case of [thyrotropin-releasing hormone](#)) to more than 200 (in the case of [follicle-stimulating hormone](#)) amino acid residues and can have molecular weights as large as 30,000. All hormones secreted by the pituitary gland are peptide hormones, as are [leptin](#) from adipocytes, [ghrelin](#) from the stomach, and [insulin](#) from the [pancreas](#).

Steroid

[Steroid hormones](#) are converted from their parent compound, [cholesterol](#). Mammalian steroid hormones can be grouped into five groups by the receptors to which they bind: [glucocorticoids](#), [mineralocorticoids](#), [androgens](#), [estrogens](#), and [progestagens](#).

History and key discoveries of endocrinology

The study of endocrinology began in China. The Chinese were isolating sex and pituitary hormones from human urine and using them for medicinal purposes by 200 BC^[4]. They used many complex methods, such as sublimation.^[5] Eventually, when [Berthold](#) noted that castrated cockerels did not develop combs and wattles or exhibit overtly male behaviour, European endocrinology began (however, it should be noted that the Chinese anticipated the science by over 1500 years.)^[6] He found that replacement of testes back into the abdominal cavity of the same bird or another castrated bird resulted in normal behavioural and morphological development, and he concluded (erroneously) that the testes secreted a substance that "conditioned" the blood that, in turn, acted on the body of the cockerel. In fact, one of two other things could have been true: that the testes modified or activated a constituent of the blood or that the testes removed an inhibitory factor from the blood. It was not proven that the testes released a substance that engenders male characteristics until it was shown that

the extract of testes could replace their function in castrated animals. Pure, crystalline [testosterone](#) was isolated in 1935.^[7]

Although most of the relevant tissues and endocrine glands had been identified by early anatomists, a more humoral approach to understanding biological function and disease was favoured by classical thinkers such as [Aristotle](#), [Hippocrates](#), [Lucretius](#), [Celsus](#), and [Galen](#), according to Freeman et al.,^[8] and these theories held sway until the advent of [germ theory](#), physiology, and organ basis of pathology in the 19th century.

In medieval [Persia](#), [Avicenna](#) (980-1037) provided a detailed account on [diabetes mellitus](#) in *The Canon of Medicine* (c. 1025), "describing the abnormal appetite and the collapse of sexual functions and he documented the sweet taste of diabetic urine." Like [Aretaeus of Cappadocia](#) before him, Avicenna recognized a primary and secondary diabetes. He also described diabetic [gangrene](#), and treated diabetes using a mixture of [lupine](#), [trigonella](#) ([fenugreek](#)), and [zedoary](#) seed, which produces a considerable reduction in the excretion of sugar, a treatment which is still prescribed in modern times. Avicenna also "described diabetes insipidus very precisely for the first time", though it was later [Johann Peter Frank](#) (1745-1821) who first differentiated between diabetes mellitus and diabetes insipidus.^[9]

In the 12th century, [Zayn al-Din al-Jurjani](#), another [Muslim physician](#), provided the first description of [Graves' disease](#) after noting the association of [goitre](#) and [exophthalmos](#) in his *Thesaurus of the Shah of Khwarazm*, the major medical dictionary of its time.^{[10][11]} Al-Jurjani also established an association between goitre and [palpitation](#).^[9] The disease was later named after Irish doctor Robert James Graves,^[12] who described a case of goiter with exophthalmos in 1835. The German [Karl Adolph von Basedow](#) also independently reported the same constellation of symptoms in 1840, while earlier reports of the disease were also published by the Italians Giuseppe Flajani and Antonio Giuseppe Testa, in 1802 and 1810 respectively,^[13] and by the English physician Caleb Hillier Parry (a friend of [Edward Jenner](#)) in the late 18th century.^[14]

In 1902 Bayliss and Starling performed an experiment in which they observed that acid instilled into the [duodenum](#) caused the [pancreas](#) to begin secretion, even after they had removed all nervous connections between the two.^[15] The same response could be produced by injecting extract of [jejunum mucosa](#) into the jugular vein, showing that some factor in the mucosa was responsible. They named this substance "[secretin](#)" and coined the term *hormone* for chemicals that act in this way.

Von Mering and Minkowski made the observation in 1889 that removing the [pancreas](#) surgically led to an increase in [blood sugar](#), followed by a coma and eventual death—symptoms of [diabetes mellitus](#). In 1922, Banting and Best realized that homogenizing the pancreas and injecting the derived extract reversed this condition.^[16] The hormone responsible, [insulin](#), was not discovered until Frederick Sanger sequenced it in 1953.

[Neurohormones](#) were first identified by [Otto Loewi](#) in 1921.^[17] He incubated a frog's heart (innervated with its [vagus nerve](#) attached) in a saline bath, and left in the solution for some time. The solution was then used to bathe a non-innervated second heart. If the vagus nerve on the first heart was stimulated, negative [inotropic](#) (beat amplitude) and [chronotropic](#) (beat rate) activity were seen in both hearts. This did not occur in either heart if the vagus nerve was stimulated. The vagus nerve was adding something to the saline solution. The effect could be blocked using atropine, a known inhibitor to heart vagal nerve stimulation. Clearly, something was being secreted by the vagus nerve and affecting the heart. The "[vagusstuff](#)" (as Loewi called it) causing

the [myotropic](#) effects was later identified to be [acetylcholine](#) and [norepinephrine](#). Loewi won the Nobel Prize for his discovery.

Recent work in endocrinology focuses on the molecular mechanisms responsible for triggering the effects of hormones. The first example of such work being done was in 1962 by [Earl Sutherland](#). Sutherland investigated whether hormones enter cells to evoke action, or stayed outside of cells. He studied norepinephrine, which acts on the liver to convert [glycogen](#) into [glucose](#) via the activation of the [phosphorylase](#) enzyme. He homogenized the liver into a membrane fraction and soluble fraction (phosphorylase is soluble), added norepinephrine to the membrane fraction, extracted its soluble products, and added them to the first soluble fraction. Phosphorylase activated, indicating that norepinephrine's target receptor was on the cell membrane, not located intracellularly. He later identified the compound as cyclic AMP ([cAMP](#)) and with his discovery created the concept of second-messenger-mediated pathways. He, like Loewi, won the Nobel Prize for his groundbreaking work in endocrinology.^{[118](#)}

Endocrinology as a profession

Although every organ system secretes and responds to hormones (including the [brain](#), [lungs](#), [heart](#), [intestine](#), [skin](#), and the [kidney](#)), the clinical specialty of endocrinology focuses primarily on the *endocrine organs*, meaning the organs whose primary function is hormone secretion. These organs include the [pituitary](#), [thyroid](#), [adrenals](#), [ovaries](#), [testes](#), and [pancreas](#).

An *endocrinologist* is a [doctor](#) who specializes in treating disorders of the endocrine system, such as [diabetes](#), [hyperthyroidism](#), and many others (see list of diseases below).

[edit] Work

The medical specialty of endocrinology involves the diagnostic evaluation of a wide variety of symptoms and variations and the long-term management of disorders of deficiency or excess of one or more hormones.

The diagnosis and treatment of endocrine diseases are guided by [laboratory](#) tests to a greater extent than for most specialties. Many diseases are investigated through *excitation/stimulation* or *inhibition/suppression* testing. This might involve injection with a stimulating agent to test the function of an endocrine organ. Blood is then sampled to assess the changes of the relevant hormones or metabolites. An endocrinologist needs extensive knowledge of [clinical chemistry](#) and [biochemistry](#) to understand the uses and limitations of the investigations.

A second important aspect of the practice of endocrinology is distinguishing human variation from disease. Atypical patterns of physical development and abnormal test results must be assessed as indicative of disease or not. [Diagnostic imaging](#) of endocrine organs may reveal incidental findings called [incidentalomas](#), which may or may not represent disease.

Endocrinology involves caring for the person as well as the disease. Most endocrine disorders are [chronic diseases](#) that need life-long care. Some of the most common endocrine diseases include [diabetes](#) mellitus, [hypothyroidism](#) and the metabolic syndrome. Care of diabetes, obesity and other chronic diseases necessitates understanding the patient at the personal and social level as well as the molecular, and the physician–patient relationship can be an important therapeutic process.

Apart from treating patients, many endocrinologists are involved in [clinical science](#) and [medical research](#), [teaching](#), and [hospital management](#).

[edit] Training

There are roughly 4,000 endocrinologists in the United States.^[*citation needed*] Endocrinologists are specialists of [internal medicine](#) or [pediatrics](#). Reproductive endocrinologists deal primarily with problems of [fertility](#) and menstrual function—often training first in obstetrics. Most qualify as an [internist](#), [pediatrician](#), or [gynecologist](#) for a few years before specializing, depending on the local training system. In the U.S. and Canada, training for board certification in internal medicine, [pediatrics](#), or [gynecology](#) after medical school is called residency. Further formal training to subspecialize in adult, [pediatric](#), or reproductive endocrinology is called a fellowship. Typical training for a North American endocrinologist involves 4 years of college, 4 years of medical school, 3 years of residency, and 3 years of fellowship. Adult endocrinologists are board certified by the [American Board of Internal Medicine](#) (ABIM) in Endocrinology, Diabetes and Metabolism.

[edit] Professional organizations

In North America the principal professional organizations of endocrinologists include The Endocrine Society,^[19] the American Association of Clinical Endocrinologists,^[20] the American Diabetes Association,^[21] the Lawson Wilkins Pediatric Endocrine Society,^[22] and the American Thyroid Association.^[23]

In the United Kingdom, the Society for Endocrinology^[24] and the British Society for Paediatric Endocrinology and Diabetes^[25] are the main professional organisations. The European Society for Paediatric Endocrinology^[26] is the largest international professional association dedicated solely to paediatric endocrinology. There are numerous similar associations around the world.

[\[edit\]](#) Patient education

Because endocrinology encompasses so many conditions and diseases, there are many organizations that provide education to patients and the public. [The Hormone Foundation](#) is the public education affiliate of [The Endocrine Society](#) and provides information on all endocrine-related conditions. Other educational organizations that focus on one or more endocrine-related conditions include the [American Diabetes Association](#), [National Osteoporosis Foundation](#), Human Growth Foundation, American Menopause Foundation, Inc., and Thyroid Foundation of America.

[\[edit\]](#) Diseases

See main article at [Endocrine diseases](#)

A disease due to a disorder of the endocrine system is often called a "hormone imbalance", but is technically known as an *endocrinopathy* or *endocrinosis*.

[\[edit\]](#) Endocrinology in popular culture

- [Dr. Lisa Cuddy](#), a character on the television show [House M.D.](#)
- [Elliot Reid](#), a character becomes an expert in the field in the [Scrubs](#) episode "[My Way Home](#)"
- [Naomi Bennett](#), a character on the television show [Private Practice](#) who is also a fertility specialist

References

- Albuzio, A., Concheri, G., Nardi, S., Dell'Agnola, G., 1994. Effect of humic fractions of different molecular size on the development of oat seedlings grown in varied nutritional conditions. In: Senesi, N., Miano, T.M. (Eds.), *Humic Substances in the Global Environment and Implications on Human Health*. Elsevier B.V., pp. 199–204.
- Arancon, N.Q., Lee, S., Edwards, C.A., Atiyeh, R.M., in press. Effects of humic acids and aqueous extracts derived from cattle, food and paper-waste vermicomposts on growth of greenhouse plants. *Pedobiologia*.
- Atiyeh, R.M., Arancon, N., Edwards, C.A., Metzger, J.D., 2000a. Influence of earthworm-processed pig manure on the growth and yield of green house tomatoes. *Bioresource Technology* 75, 175–180.
- Atiyeh, R.M., Subler, S., Edwards, C.A., Bachman, G., Metzger, J.D., Shuster, W., 2000b. Effects of vermicomposts and composts on plant growth in horticulture container media and soil. *Pedobiologia* 44, 579–590.
- Atiyeh, R.M., Edwards, C.A., Subler, S., Metzger, J.D., 2000c. Earthworm processed organic wastes as components of horticultural potting media for growing marigolds and vegetable seedlings. *Compost Science and Utilization* 8 (3), 215–223.
- Atiyeh, R.M., Dominguez, J., Subler, S., Edwards, C.A., 2000d. Changes in biochemical properties of cow manure processed by earthworms (*Eisenia andrei*) and their effects on plant-growth. *Pedobiologia* 44, 709–724.
- Atiyeh, R.M., Lee, S., Edwards, C.A., Arancon, N.Q., Metzger, J.D., 2002. The influence of humic acids derived from earthworms-processed organic wastes on plant growth. *Bioresource Technology* 84, 7–14.
- Atiyeh, R.M., Subler, S., Edwards, C.A., Metzger, J., 1999. Growth of tomato plants in horticultural potting media amended with vermicompost. *Pedobiologia* 43, 1–5.
- Barakan, F.N., Salem, S.H., Heggo, A.M., Bin-Shiha, M.A., 1995. Activities of rhizosphere microorganisms as affected by application of organic amendments in a calcareous loamy soil 2. Nitrogen transformation. *Arid Soil Research and Rehabilitation* 9 (4), 467–480.
- Bosland, P.W., Vostava, E.J., 2000. *Peppers: Vegetable and Spice Capsicums*. CABI Publishing, New York, USA.
- Bwamiki, D.P., Zake, J.Y.K., Bekunda, M.A., Woomer, P.L., Bergstrom, L., Kirchman, H., 1998. Use of coffee husks as an organic amendment to improve soil fertility in Ugandan banana production. Carbon and nitrogen dynamics in natural and agricultural tropical ecosystem 1998, 113–127.
- Cacco, G., Dell'Agnola, G., 1984. Plant growth regulator activity of soluble humic complexes. *Canadian Journal of Soil Sciences* 64, 225–228.
- Canellas, L.P., Olivares, F.L., Okorokova, A.L., Facanha, A.R., 2000. Humic acids isolated from earthworm compost enhance root elongation, lateral root emergence, and plasma H⁺-ATPase activity in maize roots. *Plant Physiology* 130, 1951–1957.
- Edwards, C.A., 1998. The use of earthworms in the breakdown and management of organic wastes. In: Edwards, C.A. (Ed.), *Earthworm Ecology*. CRC Press, Boca Raton, FL, pp. 327–354.
- Edwards, C.A., Burrows, I., 1988. The potential of earthworm composts as plant growth media. In: Edwards, C.A., Neuhauser, E. (Eds.), *Earthworms in Waste and Environmental Management*. SPB Academic Press, The Hague, Netherlands, pp. 21–32.
- Follet, R., Donahue, R., Murphy, L., 1981. *Soil and Soil Amendments*. Prentice-Hall, Inc., New Jersey.
- Grappelli, A., Galli, E., Tomati, U., 1987. Earthworm casting effect on *Agaricus bisporus* fructification. *Agrochimica* 21, 457–462.
- Johnston, A.M., Janzen, H.H., Smith, E.G., 1995. Long-term spring wheat response to summerfallow frequency and organic amendment in southern Alberta. *Canadian Journal of Plant Science* 75 (2), 347–354.
- Lee, Y.S., Bartlett, R.J., 1976. Stimulation of plant growth by humic substances. *Journal of the American Society of Soil Science* 40, 876–879.
- Maynard, A.A., 1993. Evaluating the suitability of MSW compost as a soil amendment in field-grown tomatoes. *Compost Science and Utilization* 1, 34–36.
- Muscolo, A., Felici, M., Concheri, G., Nardi, S., 1993. Effect of earthworm humic substances on esterase and peroxidase activity during growth of leaf explants of *Nicotiana plumbaginifolia*. *Biology and Fertility of Soils* 15, 127–131.
- Muscolo, A., Bovalo, F., Gionfriddo, F., Nardi, S., 1999. Earthworm humic matter produces auxin-like effects on *Daucus carota* cell growth and nitrate metabolism. *Soil Biology and Biochemistry* 31, 1303–1311.
- Mylonas, V.A., Mccants, C.B., 1980. Effects of humic and fulvic acids on growth of tobacco. I. Root initiation and elongation. *Plant and Soil* 54, 485–490.
- Nardi, S., Arnoldi, G., Dell'Agnola, G., 1988. Release of hormone-like activities from *Alloborophora rosea* and *Alloborophora caliginosa* feces. *Journal of Soil Science* 68, 563–657.
- Pascual, J.A., Garcia, C., Hernandez, T., Ayuso, M., 1997. Changes in the microbial activity of an arid soil amended with urban organic wastes. *Biology and Fertility of Soils* 24 (4), 429–434.
- SAS Institute, 2001. *SAS Procedures Guide, Version 8*. SAS Institute, Cary.
- Sims, G.K., Ellsworth, T.R., Mulvaney, R.L., 1995. Microscale determination of inorganic nitrogen in water and soil extracts. *Communications in Soil Science and Plant Analysis* 26, 303–316.
- Subler, S., Edwards, C.A., Metzger, J., 1998. Comparing vermicomposts and composts. *BioCycle* 39, 63–66.
- Tomati, U., Galli, E., 1995. Earthworms, soil fertility and plant productivity. *Acta Zoologica Fennica* 196, 11–14.
- Valdrighi, M.M., Pera, A., Agnolucci, M., Frassinetti, S., Lunardi, D., Vallini, G., 1996. Effects of compost-derived humic acids on vegetable biomass production and microbial growth within a plant (*Cichorium intybus*)-soil system: a comparative study. *Agriculture, Ecosystems and Environment* 58, 133–144.
- Wilson, D.P., Carlile, W.R., 1989. Plant growth in potting media containing worm-worked duck waste. *Acta Horticulturae* 238, 205–220.
- Zink, T.A., Allen, M.F., 1998. The effects of organic amendments on the restoration of a disturbed coastal sage scrub habitat. *Restoration-Ecology* 6 (1), 52–58.